

LICE SOLUTIONS FLOWCHART



Early lice infestation can be difficult to observe without closer inspection.

It is recommended to inspect 10 'rubbed' or light bodyweight sheep to confirm presence and degree of lousiness. Inspection should include at least 10 partings on each side of the body including the neck, withers, shoulders, rump and flank of the sheep. Adequate light is essential. Outside on a sunny day is ideal as lice don't favour light so they will wriggle making them easier to detect.

Treatment Programme
Bring shearing forward?

YES

NO

Short wool < 3 months wool growth (WG)

Long wool > 3 months (WG)

Oral

Pour-on

Ideally < 24 hours after shearing

Plunge or shower

Ideally 2 – 8 weeks after shearing

Plunge or shower

Long wool pour-on

Oral



FLEXOLT®

VANQUISH®

MAGNUM®

Water-based IGR,
safer and user-friendly

WIPE-OUT®

**ZENITH®
CONCENTRATE**

Water-based IGR,
safer and user-friendly

VANQUISH®

FLEXOLT®

Please read the label for full instructions and precautions

ACVM No's: A011971, A005997, A007704, A004558, & A006102. Schering-Plough Animal Health Ltd. Ph: 0800 800 543.
www.msd-animal-health.co.nz © 2024 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved. NZ-ZEN-240900001

COOPERS®